# Imported Sugar Beet Feed



- Dried (sugar) beet feed (unmolassed)



High digestible fibre based palatable energy feed.

## Typical Analysis (on a dry matter basis)

| Dry matter (%) | Energy<br>(MJ ME/kg<br>DM) | Crude<br>protein (%) | Oil (%) | NDF (%) | Starch (%) | Sugar (%) | DUP (%) |
|----------------|----------------------------|----------------------|---------|---------|------------|-----------|---------|
| 89.0           | 12.5                       | 9.0                  | 1.0     | 43.0    | 0.0        | 6.0       | 3.3     |

## What are you trying to achieve?

| Need                      | Feature  | Benefit  |  |
|---------------------------|--|--|--|
| Drive intake              | A highly palatable feed.                                   | Can stimulate intakes of less palatable feeds, increasing milk and meat production.                                      |  |
| Increase milk fat %       | A good source of digestible fibre.                         | Provides the building blocks for milk fat synthesis, increasing value per litre.   |  |
| Increase energy intakes   | Good levels of non-starch digestible fibre energy.         | Allows energy intakes to be increased without increasing the risk of acidosis associated with cereal feeding.            |  |
| Minimise risk of acidosis | High content of digestible fibre.                          | Assists in maintaining an optimum rumen pH.  |  |
| Feeding flexibility       | Pellet durability. Suitable for a wide range of livestock. | Designed to be suitable for use in automated and floor feeding systems. Can be transferred to feeders via auger systems. |  |

The predicted responses (benefits) assume that the specified nutrient, physical or structural dietary components are limiting livestock performance in the current ration.

## **Complementary Concentrate Feeds**

- **High starch feeds** e.g. cereals, maize meals, and confectionary and bakery products.
- **High protein feeds** e.g. soya bean meal, rapeseed meal, wheat distillers.



### Recommended daily feed rates (per head basis)



Imported Sugar Beet Feed can be fed via automated feeders, top dressed or floor fed, and used individually or as part of a blend or TMR.

| Milking Cows         | Up to 6 (typically 3)kg               |  |  |
|----------------------|---------------------------------------|--|--|
| Dry Cows             | Up to 2 kg                            |  |  |
| Replacement Heifers  | Up to 2 kg and up to 40% of the DMI   |  |  |
| Calves (to 12 weeks) | Up to 1.5 kg and up to 40% of the DMI |  |  |
| Growing Cattle       | Up to 2.5 kg and up to 40% of the DMI |  |  |
| Finishing Cattle     | Up to 5kg and up to 50% of the DMI    |  |  |
| Suckler Cows         | Up to 4 (typically 2)kg               |  |  |
| Ewes and Rams        | Up to 1 (typically 0.5) kg            |  |  |
| Hoggets and Lambs    | Up to 1 kg or up to 50% of the DMI    |  |  |

DMI = dry matter intake

## Availability, handling and storage

Unmolassed Sugar Beet Feed is usually available all year round, UK wide as bulk tipped or blown loads. Like all dry feeds, they should be stored in a secure shed, bunker, bin or hopper and kept cool, dry and free from vermin.

#### Additional information

#### Method of production

Unmolassed Sugar Beet Feed is a co-product from sugar production. Once the sugar has been diffused out from the beet, the fibrous residues are dried and then pelleted to produce the final product.

#### **Quality Assurance**

Unmolassed Sugar Beet Feed is FEMAS assured (or a recognised equivalent). Imported Sugar Beet Feed (Dried (sugar) beet feed) is listed under number 4.1.10 in the EU Catalogue of Feed Materials.

#### **Legal disclaimer**

Suggested feeding rates are produced as a guide only and many other factors may have an overriding effect on animal response; no performance guarantee can be given. Rations should be carefully balanced for energy and protein, contain sufficient forage to maintain rumen function and be fortified with an appropriate vitamin and mineral supplement. Animals must have constant access to clean water.



## **Imported Sugar Beet Feed**



## - Dried (sugar) beet feed (unmolassed)

## **Detailed Typical Analysis** (fresh basis other than where stated)

| Duri vas sitta ii | 07       | 00.0 | Calairna         | a. /l. a. | 11.00 |
|-------------------|----------|------|------------------|-----------|-------|
| Dry matter        | <u>%</u> | 89.0 | Calcium          | g/kg      | 11.00 |
| Oil A             | %        | 0.50 | Magnesium        | g/kg      | 1.30  |
| Oil B             | %        | 1.00 | Phosphorus       | g/kg      | 0.77  |
| Crude protein     | %        | 8.00 | Potassium        | g/kg      | 27.7  |
| Crude protein: DM | %        | 9.00 | Salt             | g/kg      | 1.37  |
| Fibre             | %        | 15.8 | Sodium           | g/kg      | 0.43  |
| Ash               | %        | 8.40 | Copper           | mg/kg     | 7.00  |
| ME* – in vivo     | MJ/kg DM | 12.5 | Manganese        | mg/kg     | 36.0  |
| NDF               | %        | 38.0 | Selenium         | mg/kg     | 0.09  |
| Starch            | %        | 0.00 | Zinc             | mg/kg     | 20.5  |
| Sugar             | %        | 5.50 | Saturates        | % of oil  | 23.0  |
| ERDP-FiM*         | % @ 6%   | 4.03 | Monounsaturates  | % of oil  | 11.0  |
| DUP-FiM*          | % @ 6%   | 2.90 | PUFAs            | % of oil  | 66.0  |
| DUP digestibility | %        | 70.0 | Long chain PUFAs | % of oil  | 0.00  |
| sDM               |          | 0.10 | Lysine           | % of CP   | 6.53  |
| aDM               |          | 0.20 | Methionine       | % of CP   | 1.86  |
| bDM               |          | 0.70 | Cysteine         | % of CP   | 1.63  |
| cDM               |          | 0.10 | Histidine        | % of CP   | 3.73  |
| sN                |          | 0.12 | Threonine        | % of CP   | 6.06  |
| aN                |          | 0.30 |                  |           |       |
| bN                |          | 0.65 |                  |           |       |
| cN                |          | 0.06 |                  |           |       |
|                   |          |      |                  | _         | -     |